

Relationship between the Velocity Ellipsoids of Galactic-Disk Stars and their Ages and Metallicities

V. V. Koval', V. A. Marsakov, T. V. Borkova

Institute of Physics, Southern Federal University,
Rostov-on-Don, Russia

e-mail: koval@ip.rsu.ru, marsakov@ip.rsu.ru, borkova@ip.rsu.ru

accepted 2009, Astronomy Reports, Vol. 53 No. 9, P.785-800

Abstract

Abstract The dependences of the velocity ellipsoids of F–G stars of the thin disk of the Galaxy on their ages and metallicities are analyzed based on the new version of the Geneva.Copenhagen Catalog. The age dependences of the major, middle, and minor axes of the ellipsoids, and also of the dispersion of the total residual velocity, obey power laws with indices 0.25, 0.29, 0.32, and 0.27 (with uncertainties ± 0.02). Due to the presence of thick-disk objects, the analogous indices for all nearby stars are about a factor of 1.5 larger. Attempts to explain such values are usually based on modeling relaxation processes in the Galactic disk. Elimination of stars in the most numerous moving groups from the sample slightly reduces the corresponding indices (0.22, 0.26, 0.27, and 0.24). Limiting the sample to stars within 60 pc of the Sun, so that the sample can be considered to be complete, leaves both the velocity ellipsoids and their age dependences virtually unchanged. With increasing age, the velocity ellipsoid increases in size and becomes appreciably more spherical, turns toward the direction of the Galactic center, and loses angular momentum. The shape of the velocity ellipsoid remains far from equilibrium. With increasing metallicity, the velocity ellipsoid for stars of mixed age increases in size, displays a weak tendency to become more spherical, and turns toward the direction of the Galactic center (with these changes occurring substantially more rapidly in the transition through the metallicity $[Fe/H] \approx -0.25$). Thus, the ellipsoid changes similarly to the way it does with age; however, with decreasing metallicity, the rotational velocity about the Galactic center monotonically increases, rather than decreases (!). Moreover, the power-law indices for the age dependences of the axes depend on the metallicity, and display a maximum near $[Fe/H] \approx -0.1$. The age dependences of all the velocity-ellipsoid parameters for stars with equal metallicity are roughly the same. It is proposed that the appearance of a metallicity dependence of the velocity ellipsoids for thin-disk stars is most likely due to the radial migration of stars.

Keywords: kinematics of stars, velocity ellipsoids, thin disk of the Galaxy

Introduction

The observed morphological structure of the thin disk of our Galaxy, like that of any of its subsystems, is due purely to the collected orbits of the stars making it up. Therefore, the "external" appearance of the Galaxy can be reconstructed based on the space velocities

of stars, even those located in the immediate vicinity of the Sun (assuming, of course, that this location is not distinguished in any way). Simultaneously, we can attempt to trace the dynamical evolution of the disk system using stars of various ages. A classic method for deriving such information is analysis of the age dependences of the parameters of the Schwarzschild velocity ellipsoids of stars close to the Sun. First and foremost, it is important to elucidate how the dispersions of the stellar-velocity components vary with age. The increases of all the ellipsoid axes with age can be described well by power laws of the form $\sigma_i \sim t^\gamma$, where σ_i is the dispersion of the corresponding velocity component, t the age of the stars, and γ the power-law index. It is usually supposed that these dependences arise due to relaxation processes in the Galaxy (see, for example, [1, 2]). (The alternative explanation that the velocities of the stars remain undistorted from birth, so that the age dependence of the velocity dispersion reflects variations in the dynamical state of the interstellar medium with time, is thought to be less likely.) The power-law index provides information about inhomogeneities of the gravitational potential of the Galaxy that lead to a continuous growth in the velocity dispersions of stars that are born, i. e., to "heating" of these stars. In particular, numerical modeling of heating by stochastic spiral density waves is able to explain the observed age-velocity-dispersion dependence over a broad range of power-law indices ($0.2 < \gamma < 0.7$). These numerical calculations can be used to place constraints, including constraints on parameters of the spiral structure of the Galaxy [3]. The velocities may also be perturbed by a bar in the central region of the Galaxy [4]. However, the spiral waves and the bar "operate" only in a thin layer of the Galactic disk, and are unable to explain the increase in the vertical velocity component. In this connection, molecular clouds are invoked, which can explain power-law indices in the Z direction up to values $\gamma \approx 0.26$ [5], whereas higher indices require clusters of dark matter from decaying companion galaxies disrupted by the tidal forces of our Galaxy [6].

A new era in studies of stellar kinematics arrived with the publication of the data from the HIPPARCOS satellite. Thanks to these new precision radial-velocity and proper-motion measurements, it has become clear that the velocity distribution for field stars is not uniform, with a large number of groups of stars having the same angular momenta being distinguished. These so-called moving groups or stellar flows can be arbitrarily separated into two types according to their origin. The first type of stellar flow is associated with irregularities in the Galactic potential. In particular, the Pleiades-Hyades and Sirius flows can be explained by heating of the disk by stochastic spiral waves, whose theoretical possibility was demonstrated in [3]. Another possible perturber of the stellar velocity ellipsoid is a bar at the Galactic center, whose existence is suggested by infrared observations of stars in that region [7]. It is likely that precisely a bar that generates spiral density waves [8] led to the formation of the ξ -Hercules flow (branch) at the outer Lindblad resonance [9, 10], which is located in velocity phase space at the boundary between the subsystems of the thick and thin disks of the Galaxy. Another type of flow is associated with the remnants of disrupted, fairly massive ($4 \times 10^8 M_\odot$) companions to the Milky Way [11]. Several methods for distinguishing stars in moving groups have been developed, which all lead to the identification of essentially the same stellar flows with the same members. According to numerous studies in the solar neighborhood, roughly a third of stars can be identified as members of some moving group (see, for example, [11, 12]). The stellar flows formed by inhomogeneities in the gravitational potential, i. e., by spiral waves and the bar, make up the majority. All these flows distort the velocity field of field stars of various ages, hindering the derivation of the information required to reconstruct the dynamical evolution of the Galaxy. After excluding stars in flows, the distributions for the remaining stars in $U - V$

and $V - \log t$ diagrams become smoother [13].

The relations between age and velocity dispersion for nearby field stars have already been investigated using modern astrometric and spectroscopic measurements. The power-law indices for various samples of stars lie in the range from $\gamma \approx 0.34$ [14] to $\gamma \approx 0.50$ [1, 2]. Among the most recent attempts to construct age - velocity-dispersion relations we note [15], where it was shown that "heating" of all the velocity components of Galactic-disk stars (with the mean value $\langle \gamma \rangle \approx 0.35$) has occurred over the entire lifetime of this subsystem. This conclusion concerning the continuous action of relaxation processes is supported by the results of [2], which made use of revised ages and metallicities for the stars (increasing the power-law index only slightly: $\langle \gamma \rangle \approx 0.40$). In their study of the age dependence of the vertical velocity component using data from the same catalog [15], Seabroke and Gilmore [16] concluded that a power law was not required to describe this relation, since the value of σ_W becomes saturated after ~ 4.5 billion years. Unfortunately, the relations considered in all these studies were constructed for nearby field stars, which include an appreciable number of thick-disk stars. Moreover, to obtain correct results for a sample of thin-disk stars, it is necessary to eliminate stars in moving groups. Fuller information about the kinematics of stars is provided by studies of the age dependences of all the velocity-ellipsoid parameters, not only the dispersions of the velocity components, as were considered in the papers noted above. In connection with the much discussed existence of very old and simultaneously metal-rich stars, it is also desirable to trace the variations in the velocity ellipsoids for thin-disk stars with different metallicities. An analysis of the velocity ellipsoids for thin-disk stars taking into account all these factors is the goal of the current study.

OBSERVATIONAL DATA.

The main source of data for the study was the new version of the Geneva.Copenhagen catalog, which contains the ages, metallicities, and kinematics for ≈ 14000 F-G dwarfs brighter than $V \approx 8.5^m$ [2]. (These same authors continued to refine the parameters of the catalog stars [17], but the distances, temperatures, and ages of the stars changed appreciably less than in their first revision, leaving the features of the age - velocity-dispersion relations virtually unchanged, as the authors themselves noted. The newest version of the catalog is not yet openly available.) The metallicities in the catalog were determined from the $ubvy\beta$ photometric indices corrected for interstellar reddening. The distances for approximately 75 % of the catalog stars were determined from the HIPPARCOS trigonometric parallaxes, with only parallaxes with errors less than 13 % being used. Photometric distances were used for stars without parallaxes or with parallaxes having large errors; the limiting accuracy for these is again 13 %. Photometric distances were used primarily for distant stars, while trigonometric parallaxes were used for the absolute majority of stars within 60 pc. The radial velocities in the catalog were obtained at various times as part of the CORAVEL project, and proper motions were taken from the "Tycho" catalog. The resulting claimed accuracy in the stellar velocity components is $\pm 1.5 \text{ km s}^{-1}$.

In the new catalog [2], as in the previous version [15], the most probable stellar ages are calculated using the method of [20] and the theoretical isochrones [18, 19], taking into account the uncertainties in the effective temperatures, absolute magnitudes, and metallicities. The variability of the speeds with which stars move along the evolutionary tracks was automatically taken into account. Holmberg et al. [2] found overall good agreement between their ages and the ages of [21], which were determined using the same method

and modern Yale isochrones. Allowing for the variability in the speeds of stars along their evolutionary tracks had a negative effect on stars located near turning points, where the isochrones have complex shapes, since stars evolve comparatively rapidly in such sections, and the probability of detecting a star in such stages is very low. Our tests showed that, as a rule, uncertainty in the assigned ages for stars near turning points was equal to or somewhat less than one billion years. However, since this effect was most important for comparatively young stars, metal-rich stars ended up being completely dropped from the sample, since they were taken to be located near the evolutionary stage corresponding to the depletion of the last percent of hydrogen in their cores and the beginning of core collapse (they were all assigned ages that were too young). This aspect of the method is clearly illustrated in the age-absolute magnitude diagram for single stars from the catalog with age uncertainties $\epsilon t < \pm 3$ billion years, shown in Fig. 1: the complete absence of stars at the very center of the diagram is obvious. Therefore, because of the possible unevenness in the inclusion of stars with younger ages, we must treat certain features in the age-velocity-dispersion relations with caution, especially in the age range from three to five billion years. The ages of stars with absolute magnitudes $M_V > 4.3^m$ are also uncertain, since they fall in a region on the Hertzsprung-Russell diagram where the isochrones are very dense, making the uncertainty in their ages very large. Fortunately, limiting the sample of stars based on uncertainty in the ages enables us to work with a subsample with reliable ages. Analysis shows that, in the initial catalog, a mean uncertainty $\epsilon t < \pm 3$ billion years is reached for more than 90 % of the stars, and an uncertainty of $\epsilon t < \pm 2$ billion years for 85 % of the stars. Therefore, we decided that it was reasonable to use the metallicities and ages from the catalog [2] for our statistical studies, bearing in mind the problematic range of ages.

When developing a criterion for dividing up stars according to their subsystems in the Galaxy, we took into account their chemical compositions using data from our master catalog of homogeneous spectroscopic determinations of iron and magnesium abundances [22]. Collected in this catalog are virtually all magnesium (a representative of the α elements) and iron abundances in dwarfs and subgiants of the solar neighborhood determined via synthetic fitting of high-dispersion spectra and published up to January 2004.

COMPARISON OF A REPRESENTATIVE SAMPLE OF THIN-DISK STARS.

The parameters of the velocity ellipsoids, by definition, have meaning only for nearby field stars. For stars located near the Galactic plane, the principles used to compose a particular sample of thin-disk stars could appreciably influence the resulting age dependences of the axes and other parameters of the velocity ellipsoids. It is most logical to compose a sample based on kinematic criteria, since precisely the velocities of the stars define the general structure of a subsystem. However, in this case, the velocity ellipsoids for the oldest (and consequently highest-velocity) stellar groups will depend on some criterion used to distinguish between the thin-disk and thick-disk subsystems, which will be subjective to some extent. For this reason, the index γ in the age dependences of the velocity dispersions, $\sigma_i = t^\gamma$, for the thin disk could be distorted. An alternative is to use a chemical criterion to distinguish thin-disk stars. Many studies have indicated the existence of a jump-like difference in the relative abundances of elements in thin-disk and thick-disk stars. The absolute majority of thin-disk stars have low ratios $[\alpha/Fe] < 0.25$ (see, for example, [23]). Application of this criterion enables the composition of a sample of thin-disk stars containing

no more than 10% of stars associated with the kinematic thick disk; roughly the same percent of stars that are associated with the kinematic thin disk will not be included in the sample due to their high relative magnesium abundances [24]. The membership of this small number of stars in a particular subsystem is considered uncertain [23].

Unfortunately, detailed atmospheric chemical compositions require spectra of bright objects, and so are available for only a limited number of stars. On the other hand, since the total heavy-element content (metallicity) can be derived from photometric data for much fainter stars, this quantity is known for most nearby F-G stars, making it possible to obtain statistically significant results for these stars. The metallicity can also be used as an independent criterion for distinguishing thin-disk stars. The reason is that, as was first shown in [25], the metallicity distribution for field stars displays a gap near $[Fe/H] \approx -0.5$, which naturally divides thin-disk stars from the less metal-rich stars of older subsystems of the Galaxy. As with the relative content of α elements, when this criterion is used to select thin-disk stars, just over 10% of stars associated with the kinematic thin disk are not included in the sample. Simultaneously, roughly this same fraction of thick-disk stars display high enough metallicities to be included [24]. Thin-disk stars can also be identified based on their ages, since this is a young subsystem, in which mass star formation began only ≈ 9 billion years ago, while star formation in the thick disk mostly ended ≈ 12 billion years ago [23]. However, due to the large uncertainties in the ages of individual stars, we use these ages here only in a statistical sense; i. e., purely to separate stars into groups with different ages. The above single-parameter criteria essentially select for the same stars, and the resulting samples of thin-disk stars differ only in stars that lie at the border between the two disk subsystems. In [26], we developed a kinematic criterion for distinguishing thin-disk stars, which implicitly takes into account their chemical composition. This criterion is based on our master catalog of spectroscopic derivations of the iron and magnesium contents of 867 nearby stars with accurate parallaxes [26]. We distinguish thin-disk stars simultaneously using two kinematic conditions, since either one of these on its own is insufficient. The following criterion is convenient for nearby stars: $V_{res} = (U^2 + V^2 + W^2)^{1/2} < 85 \text{ km s}^{-1}$, where U , V , W , and V_{res} are the spatial velocity components and the residual velocity of the star relative to the local centroid. The other criterion does not depend on the location of a star in its Galactic orbit, and is based on an expression proposed in [27]: $(Z_{max}^2 + 4e^2)^{1/2} < 1.05$, where Z_{max} is the maximum distance of the orbit from the Galactic plane in kpc and e is the eccentricity of the Galactic orbit. We assigned higher-velocity stars to the thick disk. We chose the limiting values of both criteria so as to minimize the number of stars with high/low relative magnesium contents in the thin/thick disk. Unfortunately, this criterion proved to be very strict, and some stars with chemical compositions more suitable for the thin disk are assigned to the thick disk.

The kinematic criterion proposed in [28] appears to be better justified from a methodological point of view. In that case, the selection of thin-disk and thick-disk stars is based on the assumption that the distributions of the stellar velocity components in both subsystems relative to the local centroid are Gaussian:

$$f(U, V, W) = \frac{1}{(2\pi)^{3/2} \sigma_u \sigma_v \sigma_w} \cdot \exp\left(-\frac{U_{LSR}^2}{2\sigma_U^2} - \frac{(V_{LSR} - V)^2}{2\sigma_V^2} - \frac{W_{LSR}^2}{2\sigma_W^2}\right).$$

Here, the ratio of the probabilities that a given star is a member of the thin disk (D) and thick disk (TD) is given by

$$D/TD = \frac{X_D}{X_{TD}} \cdot \frac{f_D}{f_{TD}}.$$

Further, stars with probabilities of belonging to each subsystem a factor of ten higher than the probability of belonging to the other subsystem are selected. The probabilities are determined using the known dispersions of each of the three spatial velocity components σ_U , σ_V , σ_W the mean rotational speed (V_{rot}) and the relative numbers of stars (X_D, X_{TD}) in both subsystems at the solar Galactocentric distance. The main disadvantages of this criterion are, first, the assumed kinematic parameters, which were derived using old data; second, the lack of self-consistency with the chemical composition of the stars; and, third, the fact that stars on the boundary between the two disk subsystems will be unidentified. As a result, this criterion proves to be even more strict than the one described above. If we reject the restrictions adopted in [28] and simply include in the thin-disk sample all stars with a higher probability of belonging to the thin disk than the thick disk, this sample includes stars whose total residual velocities reach $\sim 120 \text{ km s}^{-1}$; i.e., with kinematic properties more typical of the thick disk. In order to optimize this methodologically based kinematic criterion so that it, like our criterion described above, was simultaneously in the best possible agreement with the elemental abundances in the stars, we decided to refine the input parameters for the subsystems using the data from the catalog used. After excluding binary stars from the input catalog, we used the method of [28] to construct samples of thin-disk and thick-disk stars, then calculated the velocity-component dispersions and mean rotational velocities for each subsystem and refined the relative numbers of stars in both subsystems in the solar neighborhood. This yielded for the thin disk $\sigma_U = 31 \text{ km s}^{-1}$, $\sigma_V = 18 \text{ km s}^{-1}$, $\sigma_W = 14 \text{ km s}^{-1}$, and a speed for the asymmetric shift relative to the Sun in the direction of rotation $V_{rot} = -16 \text{ km s}^{-1}$, with these values for the thick disk being 68, 48, 49, and -53 km s^{-1} , respectively; the relative numbers of stars in these subsystems remained the same: 94 % and 6 %, respectively. The largest corrections were required for σ_V , σ_W , and (V_{rot}) for the thick-disk subsystem, which increased by 15 %-20 %. Similar values were obtained in [29] (see also [26]). We used precisely these refined parameters to select thin-disk stars from the input catalog for our study. No subsequent iterations were required, since the resulting values remained virtually constant. Figure 2a shows a plot of $V_{res} v.s. (Z_{max}^2 + 4e^2)^{1/2}$, with the thin-disk stars selected from [2] using the two criteria shown with different symbols. When the refined parameters are used, the resulting criterion becomes close to the criterion developed by us in [26]. It was not necessary to include in a given subsystem stars whose membership probabilities were a factor of ten higher than for the other subsystem; we simply selected stars whose membership probability was higher for the thin than for the thick disk. Therefore, we were able to investigate the transition between the two subsystems, obtaining undistorted results for samples not based on kinematic selection criteria. Some additional stars that were ascribed earlier to the thick disk were assigned to the thin disk. Thirty-nine such stars were found in our catalog of spectroscopic iron and magnesium abundances. Tests showed that most of these stars display low values $[Mg/Fe] < 0.25$ and high values $[Fe/H] > -0.5$, as should be observed for thin-disk stars. As a result, the kinematic criterion came into better agreement with the chemical compositions. Moreover, the resulting division of stars between the disk subsystems appears more harmonious in the diagram; therefore, we used this approach as a basis for the current study.

For our analysis, we first removed known binary stars, evolved stars ($\delta M_V > 3^m$), and stars with uncertain ages ($\epsilon t > \pm 3$ billion years) from the input catalog. Moreover, in order to eliminate boundary effects, we restricted our study to effective temperatures of 5200 – 7000 K. After applying the criterion developed above, 5965 nominally single FG thin-disk stars with reliable ages remained in the sample. (The resulting mean uncertainty

in the ages of the sample stars was $\langle t \rangle = 1.0$ billion years.)

Since our goal here is to investigate the velocity ellipsoids for thin-disk field stars and not stars in moving groups, we decided to exclude only stars of the most numerous and well known stellar flows, without applying a cumbersome mathematical procedure for this purpose. Guided by the mean velocity components of these groups determined in various studies using wavelet analyses and adaptive smoothing (see, in particular, [30, 31]), and also by the distribution of our sample of thin-disk stars in the phase space for the velocity components, we determined approximate boundaries where the density of stars appreciably exceeded the mean density in the field. The Table presents the names of the stellar flows, the ranges of velocity components relative to the Sun used to distinguish stars in the flows, and the number of stars excluded from the sample. The U , V , and W components are oriented toward the Galactic center, in the direction of the Galactic rotation, and in the direction of the North Galactic pole, respectively. The Pleiades Group was most conveniently distinguished using two velocity ranges. As a result, we eliminated about 25 % of the sample, leaving 4557 stars. The constructed UV , UW and VW distributions of the remaining sample stars demonstrate a smoother structure. Stars belonging to other known flows that were not removed from the sample are so few in number in the solar neighborhood that they are not able to significantly influence the derived parameters of the velocity ellipsoids for thin-disk field stars.

In connection with the selection of stars in the input GenevaCopenhagen catalog based purely on magnitude, the need arose to estimate the influence of selection effects due to differences in the depths of the survey for stars with different metallicities and temperatures. Hotter, higher-luminosity - i.e., younger (and therefore slower) main-sequence stars will dominate at large distances, distorting the velocity ellipsoids. Figure 2b presents the distribution of distances from the Sun for single stars from the catalog in the specified temperature range. The solid curve shows a least-squares fit to the left wing of the histogram using a power law of the form $N = \alpha R^n$. When approximating the distribution out to 60 pc, the power-law index and its 3σ errors were $n = 1.88 \pm 0.05$, as should be observed for a uniform distribution of stars in the studied volume. Therefore, for comparison, we also composed a sample out to a distance of 60 pc, which can be considered essentially complete. Only 1160 thin-disk stars remained in this sample (after excluding stars in stellar flows), which is obviously insufficient for carrying out studies for the entire volume; therefore, we will use this sample only for comparison purposes.

COMPARISON OF THE VELOCITY ELLIPSOID FOR VARIOUS SAMPLES OF THIN-DISK STARS

We calculated the parameters of the velocity ellipsoids and the velocity of the Sun relative to the local centroids using the formulas of Ogorodnikov [32], and calculated the corresponding errors using the formulas of Parenago [33]. We separated the sample into 21 age subgroups with equal numbers of stars, in order to trace variations of the velocity ellipsoids with age and clarify whether the different subsamples displayed different behavior. The first row of plots in Fig. 3 presents the age dependences of the axes $\sigma_i(t)$ for correctly selected thin-disk stars, for disk stars excluding those in moving groups, and for disk stars located within 60 pc of the Sun. (The corresponding dependences for all nearby field stars are not presented since these can be viewed in various other studies, and, in addition, the very large scatter in the points in the diagrams necessitates the use of different axis scales, hindering direct comparisons; therefore, the necessary information about these dependences will be

Table 1: Velocity components for moving-group stars relative to the Sun

Name of group	N	Velocity range of stellar streams, km s^{-1}		
		U	V	W
Sirius	367	$-12 \div 17$	$0 \div 12$	$-18 \div 6$
Hyades	555	$-43 \div -24$	$-23 \div -10$	$-17 \div -10$
Pleiades	183	$-17 \div -5$	$-31 \div -21$	$-16 \div 4$
	88	$-24 \div -10$	$-9 \div 2$	$-18 \div 6$
Coma Berenices	176	$-20 \div -6$	$-10 \div -2$	$-18 \div 3$
ξ Heracles	39	$-34 \div -27$	$-52 \div -44$	$-18 \div 4$

given in the text.) All the dependences were approximated using least-squares power laws of the form $\sigma_i \sim t^\gamma$.

We can estimate how much the behavior for the corresponding axes for stars belonging to the thin disk according to our criterion differs from that for all nearby stars (more precisely, for all single stars of the catalog [2] with age uncertainties $\epsilon t < \pm 3$ billion years). The increase in the ellipsoid axes with age can be more reliably fit with a power law for the selected thin-disk stars than for all nearby stars. The uncertainties in γ for the corresponding regression coefficients were about a factor of two smaller for all dependences for the thin-disk stars (see for comparison the age dependence of the velocity component dispersions for nearby stars in Fig. 34 of [2] and Fig. 7 of [17]). At the same time, the values of the power-law indices γ for the same dispersions were about a factor of 1.5 smaller. While the resulting indices for all nearby stars are in good agreement with the typical values obtained in other studies, $\langle \gamma \rangle = 0.42$ on average, with uncertainties ± 0.04 (compare with -0.43 ± 0.04 in [2, 17]), the indices for the correctly selected thin-disk stars were all within 3σ of $\langle \gamma \rangle = 0.28$, with uncertainties ± 0.02 (Fig. 3a). This decrease in the power-law indices is mainly due to a decrease in the values of all the axes for old subgroups, where the fraction of thick-disk stars is high. Additional analysis showed that applying stricter criteria leads to only a slight decrease in the major axes for the oldest subgroups, while the middle and minor axes remain virtually unchanged. The power-law indices for all the dependences likewise remain unchanged within the errors. (Note that, if the sample is not restricted to stars with age uncertainties $\epsilon t < \pm 3$ billion years, all the power-law indices grow, on average, by $\Delta\gamma \approx 0.03$).

Figure 3b presents the age dependences of the axes for thin-disk field stars without stars in moving groups. There are 217 stars in the subgroups, as opposed to 284 in the previous sample, which led to some increase in the uncertainties in the ellipsoid axes (see error bars in the plots); however, the scatter of the points was not increased, so that the uncertainty in the power-law indices remained the same. This figure shows that excluding stars in stellar flows increases the ellipsoid axes for the youngest subgroups beyond the errors, while leaving the axes for the old subgroups virtually unchanged. As a result, the power-law indices for all the dependences decrease somewhat, although this change remains within the errors.

Finally, Figure 3c shows the same dependences for thin-disk stars not in moving groups and located within 60 pc of the Sun. The number of stars in the subgroups is reduced to 55, leading to a large increase in the uncertainties of the axes and power-law indices (see error bars and the errors in the regression coefficients in the plots). There are no obvious shifts in these plots compared to the others: the axes and power-law indices for these dependences do not display systematic variations.

Thus, applying a correct criterion for selecting stars in the thin-disk subsystem appreciably decreased the velocity dispersions for old groups of stars, leaving these virtually unchanged for young groups. This led to an appreciable decrease in the power-law indices for the age-ellipsoid-axes relations compared to the analogous dependences for all nearby stars. Excluding stars in moving groups likewise slightly reduced the power-law indices, but this time due to an increase in the axes for young subgroups. Restricting the analysis to the most nearby stars, for which the sample of thin-disk stars in the specified temperature range is complete, did not significantly affect the results.

Note that, in all the axis plots, a break can be seen near ages of 3.5 billion years, with a sharp rise for small ages being replaced by a shallow relation for the following 2.3 billion years. This is especially clearly visible for the minor and (to a lesser extent) middle axes. (A possible origin for this effect could be the method used for the age determinations noted above.)

Let us now consider how the other parameters of the velocity ellipsoids depend on the sample used. The second row of plots in Fig. 3 presents the age dependences of the ratios of the axes for the same samples. Linear approximations of the points on all the graphs indicate small positive trends for the ratios of the minor and middle axes to the major axis, although the probability that these trends arose by chance were much higher than 5%, reflecting their small statistical significance. On all the graphs, the ratios of the middle to the major axis were < 0.60 for all ages (the larger scatter of the points about the fit curves and the larger error bars in the last graph are due primarily to the fact that the subgroups contain only a quarter as many points). Both ratios display some (though statistically insignificant) variation in the slopes of the age dependences beyond ages of $\approx 3 - 5$ billion years.

The third row of plots in Fig. 3 presents the age dependences of the vertex coordinates in Galactic coordinates for the same samples. Linear fits for all the samples display a weak decrease in the deviations in the vertex longitude (L) with increasing age (especially for the sample excluding moving groups), while the vertex latitude does not vary systematically with the age, being roughly equal to zero in all the diagrams. All these weak correlations are formally insignificant, although the constancy of the sign of the slopes for the vertex deviations in longitude for the various samples may provide evidence for their reality; it may be that these correlations would become significant for a larger sample. All the samples display distortions of the monotonicity of the age dependence of the Galactic longitude of the vertex direction near ages of 3–5 billion years (Figs. 3g, 3h, 3i).

The age dependences of the integrated parameters of the velocity ellipsoids for the components of the solar motion relative to the corresponding centroids and the Galactic coordinates of the apex of the solar motion, presented in the fourth and fifth rows of Fig. 3, are nearly identical for all three samples considered. The U and W components of the solar velocity are nearly independent of the age, while the V component increases significantly with age (as is reflected by the high correlation coefficient for this relation). Since the solar motion reflects the motion of the corresponding centroids, this indicates that, on average, the angular momentum of the stars decreases linearly with their age. The high correlation coefficients for the linear fits presented in the last row of Fig. 3 indicate a significant increase in the longitude and decrease in the latitude of the apex of the solar motion with increasing age.

VELOCITY ELLIPSOIDS FOR STARS WITH VARIOUS METALLICITIES

Analyzing the chemical composition of thin-disk stars, we found that a larger percent with anomalous relative magnesium overabundances appears with increasing metallicity [24]. This can be interpreted as evidence that these stars were formed from material that underwent a different chemical, and possibly dynamical, evolution. While stars "remember" the dynamical state of the interstellar medium from which they formed, the kinematics of stars of different metallicities can be different. In this connection, it is interesting to consider the velocity ellipsoids for stars with different metallicities. Let us first consider how the velocity ellipsoids for stars of mixed ages depend on metallicity. Since restricting the distances of the stars did not lead to any distortion of the velocity ellipsoids, we will further use our complete sample of thin-disk stars excluding stars in moving groups. To construct the diagrams in Fig. 4, the sample was divided into 11 subgroups with different metallicities, each containing 414 stars. Most of the parameters display trends with varying metallicity (see the corresponding correlation coefficients on the plots). For example, all the axes systematically decrease with increasing metallicity, with the probability that this came about by chance being $P \ll 1\%$ for all the correlations. The ratio of the middle or minor axes to the major axis, as well as the deviation of the vertex latitude also decrease with increasing metallicity, although these variations have low significance ($P \approx 20\%$), while the longitude deviations of the vertex increase strongly ($P \ll 1\%$). Some integrated parameters of the motion of the velocity ellipsoids relative to the Sun also display systematic variations with metallicity. Figure 4 shows that the correlations for the azimuthal velocity component and for both coordinates of the apex are highly significant ($P \ll 1\%$). The two other components of the solar motion are independent of metallicity. The V component displays a paradoxical behavior (first noted in [34]): the angular momentum of more metal-rich groups is lower than that of less metal-rich groups (and not higher, as could be expected!). Indeed, we saw above that the rotational velocity of stars about the Galactic center, on average, decreases linearly with increasing age, so that the angular momentum (which is proportional to the rotational velocity for nearby stars) can be taken as a statistical age indicator. On the other hand, for equal temperatures, the "isochrone" (real) ages of more metal-rich main-sequence stars are, on average, lower than those for their less metal-rich counterparts. As a result, for equal "isochrone" ages, more metal-rich stars are generally "kinematically" older than less metal-rich stars.

One feature that appears to some extent in all of the studied metallicity dependences is a break near $[Fe/H] \approx -0.25$. This break can be traced most clearly in the relations for σ_2/σ_1 , the vertex deviations (L), and the longitude of the apex; the correlation essentially disappears to the right of the break. A "leveling off" of the curves to the right of the break can also be seen in the dependences for all the ellipsoid axes.

Let us consider further the age dependences for stars in narrow metallicity ranges. For this, we divided the sample of thin-disk stars excluding stars in moving groups into four roughly equal groups separated by $[Fe/H]$ values of -0.25 , -0.13 , and 0.00 , then divided each into 12 subgroups according to age. The plots in the first row of Fig. 5 show the age dependences of the axes and dispersions of the total residual velocities for the various metallicity groups. The power-law character of the dependences is maintained in each metallicity group. Systematic variations in the power-law indices with metallicity can be traced for all the axes and the dispersion of the total residual velocity, in spite of the relatively low number of stars in the subgroups (approximately 95), which led to errors

much larger than those in Fig. 3. All the power-law indices for a given dependence display the largest values for the group with intermediate metallicity ($-0.13 < [Fe/H] < 0.00$), and systematically decrease on either side of this group (an exception is the value of γ for the minor axis of the lowest-metallicity group). The decrease in the power-law indices for the metal-poor stars is due to the increase in the velocity dispersions for the youngest subgroups, and the decrease for metal-rich stars to the decrease in the velocity dispersions for old groups. The firm detection of systematic differences in the power-law indices for stars with different metallicities will require a larger sample of thin-disk stars. A comparison of the plots in the first row also indicates that the systematic increase in the velocity dispersions with increasing metallicity for stars of mixed ages visible in Fig. 4a can be traced in stars of any age. The lower the metallicity of the group, the higher, on average, the dependences of a given type lie, with this rise increasing with age. Note that signs of the break near 3.5 billion years detected in the dependences for all thin-disk stars (Fig. 3) can be traced in all the groups. The plots in the middle row of Fig. 5 present the age dependences for the axis ratios for the same groups. These slopes do not display systematic variations with metallicity, although all the age - σ_i/σ_j correlations are positive, with the correlation coefficients differing from zero beyond the errors in some cases. More reliable determination of the slopes of these relations and differences between them will also require a larger sample of thin-disk stars. The age and metallicity dependences for the directions of the vertices of the velocity ellipsoids are presented in the third row of plots in Fig. 5. In all the plots, the deviations of the vertices in neither longitude nor latitude display significant age dependences: all the correlation coefficients are equal to zero within the errors. The complete absence of systematic variations is obvious: the corresponding correlations sometimes change sign in the transition between metallicity groups. At the same time, the systematic increase in the vertex longitude with increasing metallicity for mixed-age stars visible in Fig. 4c is characteristic for stars of all ages. Thus, the major axes of the velocity ellipsoids for stars of any age are oriented virtually toward the Galactic center for stars of low metallicity, but deviate more and more from this direction with increasing $[Fe/H]$, but only in longitude.

Let us now turn to parameters that characterize the motion of the groups as a whole rather than the dynamical state of the stars within a group. The first row of plots in Fig. 6 presents the age and metallicity dependences for the velocity components of the solar motion relative to the local centroids for the corresponding stellar groups. The negatives of these velocities represent the motion of the local centroids relative to the Sun. The components of the solar motion toward the Galactic center display a large scatter about the linear lines fit in all of the diagrams, and are independent of age (see corresponding correlation coefficients). The highly significant variations of V ($P \ll 1\%$), characterizing the angular momentum, indicate that the rotational speeds about the Galactic center for groups of any metallicity decrease with increasing age. The slopes of these dependences are approximately the same within the errors, and the paradoxical decrease in angular momentum with increasing metallicity is observed for all ages. The $V_\odot - t$ relation simply shifts upward with increasing metallicity. The velocity components directed perpendicular to the Galactic plane are roughly the same for all the groups, and do not depend on age. These properties of the velocity components are also reflected in the behavior of the coordinates of the apex of the solar motion relative to the stellar groups with different ages and metallicities. Significant trends demonstrating an increase in Galactic longitude and decrease in Galactic latitude with age are clearly visible (second row of plots in Fig. 6). The age dependence of the apex longitude is strongest for the lowest-metallicity group,

but then weakens abruptly, purely due to the increase in the apexlongitu de relative to young stellar subgroups; further, in the range $[Fe/H] > -0.25$, no systematic variations with metallicity are seen for either the slopes or the positions of the dependences. In the transition to more metal-rich groups, the shifts in latitude display a gradual decrease in their slopes in the entire metallicity range characteristic for the Galactic disk, likewise due exclusively to the decrease in the latitude of the solar-motion apex relative to young stars.

DISCUSSION

Holmberg et al. [2] constructed age - velocity-component dispersion relations for nearby stars with reliably determined ages and approximated them using power laws. The resulting power-law indices γ were 0.38, 0.38, 0.54, and 0.40 for σ_U , σ_V , σ_W , and σ_{tot} , respectively (or 0.39, 0.40, 0.53, and 0.40 in the study of Holmberg et al. [17]). Here, we have calculated the velocity-ellipsoid semi-axes, which, in general, can differ slightly from the velocity-component dispersions in a Galactic orthogonal coordinate system, since, as a rule, the major axis of the ellipsoid does not coincide with the direction toward the Galactic center. As a result, the indices for the age.axis relations for nearby stars obtained using the same data are slightly different: 0.38, 0.39, 0.48, and 0.42 for σ_1 , σ_2 , σ_3 and σ_{tot} , respectively. After excluding thick-disk stars from the sample, all the power-law indices decreased appreciably (Fig. 3a). They decreased overall by approximately a factor of 1.5 – 2 after further excluding known members of stellar flows, becoming 0.22, 0.26, 0.27, and 0.24 (Fig. 3b). The index for the velocity component perpendicular to the Galactic plane decreased most of all, although, as before, it remained the largest.

The age dependence of the W velocity component for nearby stars was considered in some detail in [16], using data from the catalog [15]. The main result of that study was to refute the results of [15, 17] concerning the continuous growth of the dispersion of W with age in the thin disk, instead finding evidence that this growth saturated ≈ 4.5 billion years ago. As a consequence, the age - σ_W relation should be fit using two power laws that intersect near 4.5 billion years. Seabroke and Gilmore [16] propose a fundamentally new mechanism for this behavior. interaction with another galaxy. Indeed, we can see a leveling off of the age - σ_3 relation near 3.5 billion years in Fig. 3c (with a similar "leveling off" also being appreciable for the other two axes). However, as we noted, this may be an artefact associated with the method used to determine the most probable ages of the stars in the input catalog. Therefore, this result must be verified using independent stellar ages derived using other methods.

Thus, we have shown that the velocity ellipsoid for thin-disk stars increases in size with increasing age, while its shape remains nearly constant (or only slightly increases). The direction of the major axis also remains nearly constant with age. the ellipsoid retains a constant, non-zero vertex deviation. At the same time, the ellipsoid's motion in the Galaxy is such that its angular momentum decreases appreciably with increasing age, while the two other velocity components remain constant. In a stationary rotating stellar system, the condition $\sigma_2/\sigma_1 = [-B/(A - B)]^{1/2}$ should be satisfied. If the Oort constants A and B are taken to be 13.7 and -12.9 $\text{km s}^{-1} \text{ kpc}$ [35], the ratio of the middle to the major axis in the Galaxy should be 0.70. Since we have found this ratio to be smaller for all ages, ($\langle \sigma_2/\sigma_1 \rangle = 0.58 \pm 0.03$), this suggests that the Galaxy should be considered to be far from stationary at any time. (Although this last conclusion can be avoided by adopting values for the Oort constants, for which the axis ratio should be equal to 0.58 (15 and -10 $\text{km s}^{-1} \text{ kpc}$)). The parameters of the velocity ellipsoids depend on metallicity. With

decreasing metallicity, the velocity ellipsoids increase in size, become more spherical, and turn in longitude toward the Galactic center. We have found an obviously paradoxical behavior for the angular momentum: for stars of equal ages, the rotational velocity about the Galactic center decreases with increasing metallicity. As a result, the mean angular momentum of the most metal-rich disk stars with ages of ≈ 2 billion years is the same as that of the least metal-rich stars with ages of ≈ 8 billion years. This cannot be explained by relaxation processes in the Galactic disk, since these cannot lead to systematic variations in the velocities of stars of a given metallicity.

CONCLUSION

Let us say a few words about the origin of kinematic differences among stellar populations in the Galactic thin disk. Our results testify that relaxation processes due to spiral density waves are able to explain the power-law character of the increase in the dispersions of the U and V stellar-velocity components in the directions toward the Galactic center and of the Galactic rotation with increasing age, with the power-law indices being $\gamma_{1,2} \approx 0.25$, and not 0.3–0.4, as has usually been thought. The somewhat higher index we have obtained for the dispersion of the vertical velocity component, $\gamma_3 \approx 0.27$, coincides within the errors with the theoretical upper limit associated with "heating" of stars by giant molecular clouds alone, without the participation of massive objects residing in them. If we suppose that relaxation processes play a determining role in the evolution of the disk, it remains unclear why the shape of the velocity ellipsoid should display only a very weak time dependence, while simultaneously remaining far from equilibrium. On the other hand, the significant decrease in the mean angular momentum of stars with increasing age testifies that the stars "remember" the dynamical state of the interstellar medium from which they formed, and that relaxation processes have not fully distorted the initial stellar-velocity distribution. The existence of metallicity dependences for the velocity ellipsoids for stars of the same age is also difficult to explain purely using relaxation processes, although corresponding age dependences for different metallicity groups differ little. The larger radial and azimuthal velocity dispersions and nearly zero vertex deviations of stars with $[Fe/H] < -0.25$ (especially young ones) provide evidence that they formed from interstellar gas containing a substantial fraction of matter that had fallen in from outer parts of the Galaxy. Support for this hypothesis comes from their enhanced abundances of α elements, compared to the Sun [24]. However, the high angular momentum carried by young, metal-poor stars and their small (as for other young disk stars) velocity dispersions perpendicular to the Galactic disk are not fully in agreement with this picture. It is also difficult to understand the linear growth in the vertex deviation, which is the same for stars of any age, and the paradoxical decrease in angular momentum with increasing metallicity. Another problem that may be related to these is the sharp decrease in the power-law indices for the age dependences of the velocity dispersions for stars with $[Fe/H] > 0.0$. One of the most promising ways to explain all these contradictions may be a detailed analysis of radial migrations of stars of various metallicities, which have ended up near the Sun at the current time as a result of the action of inhomogeneities in the Galactic potential. The analysis of the orbits of nearby stars carried out in [36] showed that more than half of stars with $[Fe/H] < -0.4$ have orbits whose properties suggest they were born in the outer regions of the Galactic disk, while most stars with $[Fe/H] > 0.1$ were born in inner regions.

Thus, our use of the largest sample of nearby field stars with accurately determined

physical and kinematic parameters currently available has enabled us to identify several important dependences, that can ultimately provide a better understanding of the evolution of the thin disk. Nevertheless, as we have noted in the text, we were not able to draw statistically significant conclusions in some cases.

References

- [1] B. Fuchs, C. Dettbarn, H. Jahreiss, and R. Wielen, in Dynamics of Star Clusters and the Milky Way, Ed. by S. Deiters, B. Fuchs, A. Just, R. Spurzem, and R. Wielen, ASP Conf. Ser. **228**, 235 (2001)
- [2] J. Holmberg, B. Nordstrom, and J. Andersen, *Astron. Astrophys.* **475**, 519 (2007)
- [3] R.S. De Simone, X. Wu, and S. Tremaine, *Mon. Not. R. Astron. Soc.* **350**, 627 (2004)
- [4] D. Chakrabarty, *Astron. Astrophys.* **467**, 145 (2008)
- [5] J. Hanninen and C. Flynn, *Mon. Not. R. Astron. Soc.* **347**, 732 (2002)
- [6] A.J. Benson, C.G. Lacey, C.S. Frenk, et al., *Mon. Not. R. Astron. Soc.* **351**, 1215 (2004)
- [7] C. Babysiaux and G. Gilmore, *Mon. Not. R. Astron. Soc.* **358**, 1319 (2005)
- [8] R. Fux, *Astron. Astrophys.* **373**, 511 (2001)
- [9] W. Dehnen, *Astrophys. J. Lett.* **524**, L35 (1999)
- [10] W. Dehnen, *Astron. J.* **119**, 800 (2000)
- [11] A. Helmi, J.F. Navarro, B. Nordstrom, et al., *Mon. Not. R. Astron. Soc.* **365**, 1319 (2006)
- [12] B. Famaey, A. Jorissen, X. Luri, et al., *Astron. Astrophys.* **431**, 165 (2005)
- [13] R. Assian, F. Figueras, J. Torra, and B. Chen, *Astron. Astrophys.* **351**, 427 (1999)
- [14] J. Binney, W. Dehnen, and G. Bertelli, *Mon. Not. R. Astron. Soc.* **328**, 658 (2000)
- [15] B. Nordstrom, M. Mayor, J. Andersen, et al., *Astron. Astrophys.* **418**, 989 (2004)
- [16] G.M. Seabroke and G. Gilmore, *Mon. Not. R. Astron. Soc.* **380**, 1358 (2007)
- [17] J. Holmberg, B. Nordstrom, and J. Andersen, *Astron. Astrophys.* (2009, in press); arXiv:0811.3982v1 [astro-ph] (2008).
- [18] L. Girardi, A. Bressan, C. Chiosi, et al., *Astron. Astrophys. Suppl. Ser.* **141**, 371 (2000)
- [19] B. Salasnich, L. Girardi, A. Weiss and C. Chiosi, *Astron. Astrophys.* **361**, 1023 (2000)
- [20] B.R. Jorgensen and L. Lingren, *Astron. Astrophys.* **436**, 127 (2005)
- [21] Y. Takeda, *Publ. Astron. Soc. Japan* **59**, 345 (2007)
- [22] T.V. Borkova and V.A. Marsakov, *Astron. Zh.* **82**, 453 (2005) [*Astron. Rep.* **49**, 405 (2005)]

- [23] K. Fuhrmann, in *The First Stars*, Proc. MPA/ESO Workshop, Ed. by A. Weiss, T.G. Abel, and V. Hill (Springer, 2000), p. 68.
- [24] V.A. Marsakov and T.V. Borkova, *PismaAstron. Zh.* **32**, 419 (2006) [*Astron. Lett.* **32**, 376 (2006)]
- [25] V.A. Marsakov and A.A. Suchkov, *Astron. Zh.* **54**, 1233 (1977) [*Sov. Astron.* 21, 700 (1977)]
- [26] V.A. Marsakov and T.V. Borkova, *PismaAstron. Zh.* **31**, 577 (2005) [*Astron. Lett.* 31, 515 (2005)]
- [27] R.G. Gratton, E. Carretta, R. Claudi, et al., *Astron. Astrophys.* **404**, 187 (2003)
- [28] T. Bensby, S. Feldzing, and I. Lundstrom, *Astron. Astrophys.* **410**, 527 (2003)
- [29] G. Gilmore and R.F.G. Wyse, in *The Galaxy*, Ed. by G. Gilmore and B. Carswell, NATO ASI Ser. C (Reidel, Dordrecht, 1987), v. 207, p. 247.
- [30] V.V. Bobylev and A.T. Ba@ikova, *Astron. Zh.* **84**, 418 (2007) [*Astron. Rep.* 51, 372 (2007)]
- [31] T. Antoja, F. Figueras, D. Fernandez, and J. Torra, *Astron. Astrophys.* **490**, 135 (2008)
- [32] K.F. Ogorodnikov, *Dynamics of Stellar Systems* (Izd. Fiz.-Mat. Liter., Moscow, 1958), p. 84 [in Russian].
- [33] P.P. Parenago, *Tr. Gos. Astron. Inst.* **20**, 26 (1951)
- [34] V.A. Marsakov, A.A. Suchkov, and Yu.G. Shevelev, *Astrophys. Space Sci.* **172**, 51 (1990)
- [35] V.V. Bobylev, *Pisma Astron. Zh.* **30**, 185 (2004) [*Astron. Lett.* 30, 159 (2004)]
- [36] M. Haywood, *Mon. Not. R. Astron. Soc.* **388**, 1175 (2008)




Figure 1: Ageabsolute magnitude diagram for single stars of the catalog [2] with age uncertainties $\epsilon t < \pm 3$ billion years.




Figure 2: (a) Relation between $\sqrt{Z_{\max}^2 + 4e^2}$ and the residual velocities relative to the local centroid for stars with age uncertainties $\epsilon t < \pm 3$ Gyr and (b) the distribution of distances from the Sun for single stars from the catalog [2]. Hollow circles show thin-disk stars selected according to the criterion of [26], and filled circles stars selected according to the new criterion. The pluses show all remaining stars of the catalog. The curve is a power-law fit to the distribution within 60 pc from the Sun.




Figure 3: Age dependences of the velocity ellipsoids for all single F-G thin-disk stars with age uncertainties $\epsilon t < \pm 3$ Gyr. The first column shows plots for stars selected according to our criterion, the second excludes stars of moving groups, and the third excludes stars more than 60 pc from the Sun. The age dependences for the axes of the velocity ellipsoids and the total residual velocity (upper row), ratios of the ellipsoid axes (second row), vertex coordinates (third row), solar-velocity components relative to the corresponding centroids (fourth row), and apex coordinates for the solar motion (fifth row) are presented. The curves in the upper row of plots are power-law fits, the titles above the plots give the power-law indices with their errors, and the bars show the errors in the




Figure 4: Metallicity dependences of the (a) velocity-ellipsoid axes and the total residual velocity, (b) the ratios of the ellipsoid axes, (c) the coordinates of the vertex, (d) the components of the solar velocity relative to the corresponding centroids, and (e) the coordinates of the apex of the solar motion. The solid lines are linear fits and the numbers indicate the corresponding correlation coefficients. The mean error bars for the parameters are less than the size of the symbols in all the figure panels.




Figure 5: Age dependences of the velocity ellipsoids for thin-disk stars for four metallicity groups. The metallicity ranges are indicated above. The notation is as in Fig. 3.




Figure 6: Age dependences of the solar velocity components relative to the corresponding centroids (first row) and the coordinates of the solar-motion apex (second row) for four metallicity groups. The metallicity ranges are indicated above. The notation is as in Fig. 3.